

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SLUGGIT ***PRIMA 30***

SLUG AND SNAIL BAIT

SLUGGIT
PRIMA 30

NET CONTENTS:
25kg

ACTIVE CONSTITUENT: 30 g/kg METALDEHYDE



CONTROLS
SLUGS &
SNAILS IN
AGRICULTURAL
AND
HORTICULTURAL
CROPS AND
DURING PASTURE
ESTABLISHMENT

**FOR THE CONTROL OF SLUGS & SNAILS IN
AGRICULTURAL AND HORTICULTURAL
CROPS AND DURING PASTURE ESTABLISHMENT**

NET CONTENTS:
25kg

APVMA Approval Number 91239/131484

GrowthCare
Your growth. Our care.

BATCH NUMBER

MANUFACTURING DATE

**SLUGGIT
PRIMA 30**

NET CONTENTS:
25kg

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply to crops once windrowed.

DO NOT apply to edible plant parts or to crops where bait may be collected with harvested commodity.

DO NOT place baits in heaps. Clean up spilled baits immediately, including after filling applicator.

CROP/SITUATION	PEST	RATE	CRITICAL COMMENTS
Agricultural crops Cereals, oilseeds (including poppies) pulses	Slugs and snails	3 kg/ha or 30 g/100 m ²	Broadcast evenly over the surface of the ground where slugs and snails are active.
Horticultural crops Fruit (including grapevines) & vegetables		6 kg/ha or 60 g/100 m ²	
Pasture establishment*		3 to 6 kg/ha	* Pasture crops only: SLUGGIT PRIMA 30 bait is to be used only during the establishment stages of pastures. Broadcast pellets evenly over the crop area to be protected either prior to planting, post-plant pre-emergence or early post-emergence (prior to 4 leaf stage). Use higher rate where there are large populations of slugs or snails and where wet and cool climatic conditions favour slug and snail movement.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

This product contains denatonium benzoate deterrent to help with pet safety.

WITHHOLDING PERIODS

HARVEST

HORTICULTURAL CROPS, VINEYARDS: DO NOT APPLY LATER THAN 7 DAYS (1 WEEK) BEFORE HARVEST.

AGRICULTURAL CROPS: DO NOT APPLY LATER THAN 8 WEEKS BEFORE HARVEST.

GRAZING

PASTURES, PULSES, OILSEEDS (INCLUDING POPPIES), WINTER CEREALS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.

ALL OTHER CROPS: DO NOT ALLOW STOCK OR POULTRY TO GRAZE TREATED CROPS.

GENERAL INSTRUCTIONS

SLUGGIT PRIMA 30 bait has a stomach and contact action in slugs and snails that induces heavy secretion of mucous with the consequence that slugs and snails die in a reasonably short time from dehydration. SLUGGIT PRIMA 30 bait is particularly active against molluscs. If the product is applied according to the directions for use, non-target organisms will not be affected. Addition of a bittering agent has made SLUGGIT PRIMA 30 bait distasteful to mammals and birds. Best results are achieved by application after rain, watering or during damp and humid weather when the slugs and snails are most actively seeking food but not when rain is imminent. Scatter bait evenly using an appropriate applicator by ground or air, or direct drill with seed for slug control. Clean up spilled bait after filling applicator or whenever the bait is spilt.

PRECAUTIONS

DO NOT use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the product or used containers.

PROTECTION OF DOMESTIC ANIMALS AND LIVESTOCK

Keep away from animals, poultry, ducks and domestic pets as accidental poisoning may be fatal. DO NOT graze or cut for stockfeed any area while SLUGGIT PRIMA 30 bait is visible in the plant foliage. If pets or livestock are poisoned, contact a veterinary surgeon.

STORAGE AND DISPOSAL

Store in the closed original container in a dry, cool, well ventilated area out of direct sunlight. **STORE IN A LOCKED ROOM OR PLACE OUT OF THE REACH OF CHILDREN, ANIMALS, FOOD, FEEDSTUFF, SEED AND FERTILISERS.** Shake and empty contents into application equipment. Do not dispose of unused product on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

When opening the container and loading and if applying by hand, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length rubber gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 1800 764 766.

MANUFACTURED BY

Orchard Agrikem (Pty) Ltd
PO Box 94, Worcester, 6849
Western Cape, South Africa
www.orchardagri.co.za

DISTRIBUTED BY

Growthcare (Pty) Ltd
8 Hasler Road
Osborne Park, 6017 WA
Australia
email: info@growthcare.com.au

For more information, visit www.growthcare.com.au

SLUGGIT IS A REGISTERED TRADEMARK.

**CONTROLS
SLUGS &
SNAILS IN
AGRICULTURAL
AND
HORTICULTURAL
CROPS AND
DURING PASTURE
ESTABLISHMENT**